Coredem meeting – Barcelona, November 19th 2021







Is your gaze your aim?

Eye position in reward gambling and the role of orbito-frontal cortex in encoding the value of visually cued offers


<u>Demetrio Ferro</u>^{1,2,*}, Anna Rifé Mata^{1,2}, Tyler Cash-Padgett³, Maya Zhe Wang³, Benjamin Hayden³, Rubén Moreno Bote^{1,2}

¹Center for Brain and Cognition (CBC), Universitat Pompeu Fabra (UPF), 08002, Barcelona – ES;


²Department of Information and Communication Technologies, Universitat Pompeu Fabra (UPF), 08002, Barcelona – ES;

³Dept. of Neuroscience, Center for Magnetic Resonance Research, Center for Neuroeng., University of Minnesota, MN55455, Minneapolis – USA;


*demetrio.ferro@upf.edu





Fixate




acquire fixation at center of the screen










Delay 1




blank screen









Delay 2




blank screen



Re-fixate




re-acquire fixation at center of the screen



Choice-go




saccade to chosen offer side



Choice-made



hold chosen offer side for at least +200ms



Reward





reward is provided





Motivations


- Is the gaze position relevant for the reward gambling task execution?
- Can we use the gaze position as a marker of what is the animal mentally picturing during task execution in, particular during delay times?














Trials pooled with first offer re-referenced to Left side.

Left is best, E(Left)>E(Right)Right is best, E(Right)>E(Left)







int



Generalized Linear Model (GLM) for behavioral choice



Neural Data




Carmichael, S.T., and Price, J.L. (1994). Architectonic subdivision of the orbital and medial prefrontal cortex in the macaque monkey. J. Comp. Neurol.346,366–402.

Subject 1

area	session	#cells	# trials
BA13	12/07/17	51	643
BA13	12/08/17	59	700
BA11	12/09/17	24	697
BA11	12/10/17	29	603
	Total	163	2643

Subject 2

area	session	#cells	# trial
BA11	3/06/19	18	1015
BA11	3/07/19	32	323
BA11	3/08/19	9	1084
BA11	3/11/19	26	906
	total	85	3328



- 2 Subjects
- 8 Sessions
- 248 Cells

Data acquisition




Tyler Cash-Padgett, Maya Zhe Wang, Benjamin Hayden, Hayden Lab, Dept. of Neuroscience, Center for Magnetic Resonance Research, Center for Neuroengineering, University of Minnesota, Minneapolis, USA;


Two adult male rhesus macaques (macaca mulatta) served as experimental subjects. All procedures were approved by the University Committee on Animal Resources at the University of Rochester and at the University of Minnesota, conducted in compliance with the Public Health Service's Guide for the Care and Use of the Animals.

Motivations

- Are task-relevant variables encoded by OFC cells?
- Is the gaze position relevant in the neural process of encoding the offer values?

GLM for OFC spiking activity






Hypothesis:

Focusing on the Left offer EV, E(L)

if the monkey looks at Left/Right side, is the **E(L)** coding in OFC affected?





Test:


Consider trials where monkey mostly


- LookL: tR/(tR+tL)<0.5
- LookR: tR/(tR+tL)>0.5
- GLM for E(L): Look Left $\eta \approx \text{Poiss}(f^{-1}(\beta_{0,L} + \beta_L \cdot E(L)))$
- GLM for E(L): Look Right $\eta \approx \text{Poiss}(f^{-1}(\beta_{0,R} + \beta_R \cdot \mathbf{E}(\mathbf{L})))$

 β_L vs β_R ??













GLM for **E(L)**: LookL vs LookR






GLM for **E(R)**: LookL vs LookR






Conclusions


• The gaze position has a significant role in the reward gambling task execution: the fraction of time spent at either screen side is predictive of the chosen side;



• Task-relevant variables are encoded by a significant fraction of OFC cells, including the fraction of time spent inspecting either screen side;



• The gaze position is relevant in the process of encoding offer values: looking at either side possibly yields stronger coding of the ipsi-later offer EV.







TCN upf.edu/web/tcn Research Group on Theoretical and Cognitive Neuroscience

TCN Lab

Rubén Moreno Bote Anna Rifé Mata Alice Vidal Jorge Ramirez Ruiz Chiara Mastrogiuseppe Carolina Schneider Bender Devin Ozbağcı Dmytro Grytskyy Farhad Razi Sofia Lawrie Alireza Valyan Francesco Damiani Fatma Aboalasaad

Collaborators

Benjamin Hayden, Tyler Cash-Padgett, Maya Zhe-Wang,

Hayden Lab, <u>haydenlab.com</u> University of Minnesota, Dept. Neuroengineering, Minneapolis, USA.


Fundings






Thank you for your attention.


Eye Data





- → Subject is inspecting the CURRENT offer (on screen)
- → Subject is not interested in CURRENT offer (on screen) since it is not the best
- → Subject drifts the gaze to NEXT offer location (blank screen) for a better value
- → Subject drifts the gaze back to PREVIOUS location (blank screen) for a better value
- → Subject correctly holds the gaze to choose the BEST offer


Eye Data



Experimental paradigm

Reward gambling task

Orbito-Frontal Cortex (OFC)




Data acquisition




Tyler Cash-Padgett, Maya Zhe Wang, Benjamin Hayden, Hayden Lab, Dept. of Neuroscience, Center for Magnetic Resonance Research, Center for Neuroengineering, University of Minnesota, Minneapolis, USA;


Two adult male rhesus macaques (*macaca mulatta*) served as experimental subjects. All procedures were approved by the University Committee on Animal Resources at the University of Rochester and at the University of Minnesota, conducted in compliance with the Public Health Service's Guide for the Care and Use of the Animals.




GLM for neural spiking activity




x₁, w₁, Left offer EV x₂, w₂, Right offer EV x₃, w₃, Left offer VAR x₄, w₄, Right offer VAR x₅, w₅, order 1stL=1 x₆, w₆, tR/(tR+tL) n.s. P<0.05




GLM for neural spiking activity




x₁, w₁, Left offer EV x₂, w₂, Right offer EV x₃, w₃, Left offer VAR x₄, w₄, Right offer VAR x₅, w₅, order 1stL=1 x₆, w₆, tR/(tR+tL) n.s. P<0.05




GLM for neural spiking activity



x₁, w₁, Left offer EV x₂, w₂, Right offer EV x₃, w₃, Left offer VAR x₄, w₄, Right offer VAR x₅, w₅, order 1stL=1 x₆, w₆, tR/(tR+tL)





 β_R

 $\boldsymbol{\beta_L}$

 β_L β_R

GLM for **E(L)**:

LookL vs LookR


GLM for **E(R)**:


LookL vs LookR


GLM for


E(R) - E(L):

LookL vs LookR










What if we used more bins for Look L vs Look R? i.e. tR/(tR+tL) binned as [0, 0.25, 0.5, 1]



What if we used more bins for Look L vs Look R? i.e. tR/(tR+tL) binned as [0, 1/6, 2/6, 3/6, 4/6, 5/6, 1]

